Contents Volume 88, 1991

Coral reef geomorphology as a function of seasonal prevailing currents and larval dispersal
P. W. Sammarco, J. C. Andrews (Townsville, Qld., Australia) and M. J. Risk (Hamilton, Ont., Canada) 1
A continuous record of Holocene pollen from Harris Lake, southwestern Saskatchewan, Canada
M. A. Sauchyn and D. J. Sauchyn (Regina, SK, Canada) 13
Palaeobathymetry and palaeogeography of the Bakony Eocene Basin in western Hungary
M. Báldi-Beke and T. Báldi (Budapest, Hungary) 25
A Late Pleistocene "initial" vegetation at Vrøgum, west Jutland (Denmark)
E. Kolstrup (Sønderborg, Denmark) 53
Paleobiogeographic relationships of Andean angiosperms of Cretaceous to Pliocene age
D. W. Taylor (New Haven, CT, USA) 69
Integrated vertebrate, invertebrate and plant taphonomy of the Fossil Forest area (Fruitland and Kirtland formations: Late Cretaceous), San Juan County, New Mexico, USA
A. P. Hunt (Albuquerque, NM, USA) 85
Dispersion et peuplement de *Soudanella laciniosa* Apostolescu, 1961 (Ostracoda), au Danien/Paléocène supérieur durant l'ouverture de l'Atlantique équatorial [Dispersal of *Soudanella laciniosa* Apostolescu 1961 (Ostracoda) Danian to upper Paleocene during the opening of the equatorial Atlantic Ocean.]
G. Carbonnel (Villeurbanne, France) 109
Trace element variations and dating of a Late Pleistocene Tasmanian speleothem
A. Goede (Hobart, Tasmania, Australia) and J. C. Vogel (Pretoria, South Africa) 121
Les crises du Dévonien supérieur par l'étude des faunes d'ostracodes dans leur cadre paléogéographique [The Upper Devonian crises with reference to ostracod faunas in their paleogeographic framework.]
F. Lethiers and D. Raymond (Paris, France) 133
The mid-Cretaceous palaeoenvironment of central Southern Africa (Orapa, Botswana)
R. J. Rayner, S. B. Waters, I. J. McKay (Wits, South Africa), P. N. Dobbs (Johannesburg, South Africa) and A. L. Shaw (Lobatse, Botswana) 147
Prey bone utilization by predatory dinosaurs
A. R. Fiorillo (Pittsburgh, PA, USA) 157
Trace-fossil tiering in flysch sediments: examples from the Guipúzcoan flysch (Cretaceous–Paleogene), northern Spain
S. Leszczyński (Krakow, Poland) 167
Palaeoecological significance of *Lithophaga* borings in Albian stromatolites, SE Nigeria
E. B. Akpan (Calabar, Nigeria) 185
Mesozoic Tethyan strata of Thakkhola, Nepal: evidence for the drift and breakup of Gondwana
F. M. Gradstein (Dartmouth, N.S., Canada), M. R. Gibling (Halifax, N.S., Canada), M. Sarti (Castigliane Cosentino Scalo, Italy), U. von Rad (Hannover, Germany), J. W. Thurow (Bochum, Germany), J. G. Ogg (West Lafayette, IN, USA), L. F. Jansa (Dartmouth, N.S., Canada), M. A. Kaminski (Kiel, Germany) and G. E. G. Westermann (Hamilton, Ont., Canada) ... 193
Distributions of halloysite–metahalloysite and gibbsite in tropical mountain paleosols: relationship to Quaternary paleoclimate
W. C. Mahaney (North York, Ont., Canada) 219
Use of organic carbon content as a chronostratigraphic tool
B. Carpentier, J. P. Gely, A. Y. Huc, B. Durand (Rueil-Malmaison, France), M. M. Blanc-Valleron (Paris, France) and M. Ansart (Mulhouse, France) 231
Paleoclimatology and paleobotany of the Eocene Chumstick Formation, Cascade Range, Washington (USA): a rapidly subsiding alluvial basin
J. E. Evans (Bowling Green, Ohio, USA) 239
Les échinides Hemiasteridae: marqueurs écologiques de la plate-forme castillane et navarro-cantabre (Espagne) au Crétacé supérieur [Hemiasterid echinoid ecological zonations of the Late Cretaceous Castillian and Navarro-Cantabrian platforms (Spain)]
D. Néraudeau and M. Floquet (Dijon, France) 265